

Mortality from 1 to 16–18 years in bilateral cerebral palsy

Gillian Baird,¹ Elizabeth Allen,² David Scrutton,^{1,3} Adrienne Knight,⁴ Anne McNee,¹ Elspeth Will,¹ Diana Elbourne²

¹Newcomen Centre, Guy's and St Thomas' NHS Trust, London, UK

²Medical Statistics Unit, London School of Hygiene and Tropical Medicine, London, UK

³Neurosciences Unit, Institute of Child Health, University College, London, UK

⁴Department of Paediatrics, King's College London, London, UK

Correspondence to

Professor G Baird, Newcomen Child Development Centre, Guy's and St Thomas' NHS Trust, London SE1 9RT, UK; gillian.baird@gstt.nhs.uk

Accepted 22 April 2010
Published Online First
15 June 2010

ABSTRACT

Objective To ascertain mortality from 1 to 18 years, and predictors of mortality.

Design Long-term follow-up of population cohort born 1989–1992.

Setting Births in South East Thames Region.

Patients 346 children with bilateral cerebral palsy (CP).

Interventions Not applicable.

Main outcome measures Mortality; predictors of mortality.

Results 98% of the cohort were traced. 61/340 (17.9%) had died by age 16–18 years at a steady mortality. The main predictive factor was severity of impairment of functional ability (hazard ratio 5.7, 95% CI 2.1 to 15.0 for poor hand manipulation; 6.8 (1.9 to 23.9) for severe communication problems).

Conclusions Although there were deaths throughout the childhood and teenage years, the majority of children with bilateral CP are likely to survive to adulthood, especially if they do not have major functional impairment at 2 years. This confirms findings of other studies of children with CP.

BACKGROUND

Mortality in children with cerebral palsy (CP) is greatly increased compared with unaffected children.^{1–3} Cohort studies suggest that severity of functional disability is the best predictor of mortality. They are based on CP registers and multiple clinicians' assessments, variably include postneonatal causes and include children with unilateral CP, whose brain involvement may have a different functional impact from that of bilateral CP.

We therefore report the follow-up to age 16–18 years of a cohort of children with bilateral CP alive at 12 months of age whose physical examination was made uniquely by one clinician (DS).

METHODS

The methods have been previously described.⁴ Briefly, all children with movement impairment born to mothers resident in a geographically defined UK region in 1989–1992 were physically examined by DS, and those with bilateral CP were followed up. Intellectual and communicative ability was directly assessed by GB in approximately half the children and information obtained from local child development services for all. As the original study focused on the development of hips which lacked the influence of previous walking experience, only children diagnosed with CP by 15 months were included. Children who died before 12 months of age were excluded.

What is already known on this topic

- ▶ Most children with cerebral palsy (CP) live to adulthood.
- ▶ Severity of functional disability is the best predictor of mortality.

What this study adds

- ▶ Even among the category of children with bilateral CP most live to adulthood, but there is an 18% mortality from those alive at 1 year with deaths occurring steadily to 16–18 years.
- ▶ Based on factors known at age 2 years, the main independent predictor of mortality is the severity of motor impairment affecting communication and manual function.

Between 2006 and 2008 the original cohort was followed up until death or censored at the latest date they were known to be alive. Death certificates were obtained from the Office for National Statistics. Factors (shown in table 1) known by age 2 years (± 6 months) were used for the prediction of death. Children dying before 18 months were excluded.

Statistical analysis

Cox proportional hazard models were used to examine the association between survival and explanatory factors. In all initial analyses, the five-point categorical scales and the categorised birthweight variable were treated as continuous measures; this approach was supported by tests for linear trend.

Following univariate analyses, multivariate analyses were carried out to consider the effect of several disabilities. In order to select the variables for the multivariate analysis, a full model on all explanatory variables was fitted, the least significant term was removed and the model re-estimated. This was repeated until all remaining variables were significant at the 5% level.

In order to allow comparison with other studies, we examined whether there was any difference in survival between early and late onset cases. Two approaches were used: first, any acquired cause up to 28 days postnatally (as used by Hutton *et al*)¹ and, second, an acquired cause with onset later

Short report

Table 1 Univariate analysis of mortality predictors

	Alive (n=279)	Dead (n=57)	Hazard ratio (95% CI)
	No (%)	No (%)	
Early onset (insult before 28 days after birth)	258 (92.5)	50 (87.7)	
Late onset (onset 28 days to 15 months)	21 (7.5)	7 (12.3)	1.6 (0.7 to 3.6)
Any postnatal cause			
No	228 (81.7)	45 (79.0)	
Yes	51 (18.3)	12 (21.1)	1.2 (0.6 to 2.3)
Gender			
Male	170 (60.9)	28 (49.1)	
Female	109 (39.1)	29 (50.9)	0.7 (0.4 to 1.1)
Birth weight (g)			
≤1500	62 (22.2)	5 (8.8)	
1500–1999	48 (17.2)	7 (12.3)	
2000–2499	28 (10.0)	7 (12.3)	
>2500	140 (50.2)	38 (66.7)	1.4 (1.1 to 1.8)**
Missing	1 (0.4)	0	
Gestational age (weeks)			
Mean (SD)	34.9 (5.4)	37.3 (4.7)	1.1 (1.0 to 1.2)**
Missing	1	0	
Multiple births			
1	247 (88.5)	55 (96.5)	
2	17 (6.1)	1 (1.8)	
3	10 (3.6)	1 (1.8)	
4	2 (0.7)	0	0.5 (0.2 to 1.3)
Missing	3 (1.1)	0	
Severity of non-motor functional disability			
Communication problems			
Mean (SD)	1.6 (1.3)	3.5 (0.9)	4.4 (3.1 to 6.4)***
No (score 0, 1 or 2)	195 (69.9)	4 (7.0)	
Yes (score 3 or 4) (severe and profound)	81 (29.0)	53 (93.0)	
Missing	3 (1.1)	0	
Severe visual impairment (best eye)			
Mean (SD)	0.5 (1.1)	2.0 (1.7)	1.8 (1.5 to 2.1)
No (score 0, 1 or 2)	247 (88.5)	28 (49.1)	
Yes (score 3 or 4) (<6/60)	28 (10.0)	26 (45.6)	
Missing	4 (1.4)	3 (5.3)	
Severe hearing impairment (best ear)			
Mean (SD)	0.1 (0.5)	0.2 (0.6)	1.1 (0.7 to 1.7)
No (score 0, 1 or 2)	271 (97.1)	49 (86.0)	
Yes (score 3 or 4) (<70 dB aided)	5 (1.8)	1 (1.8)	
Missing	3 (1.1)	7 (12.3)	
Hydrocephalus			
No	259 (92.8)	53 (93.0)	
Yes	20 (7.2)	4 (7.0)	1.0 (0.4 to 2.8)
Shunt			
No	263 (94.3)	53 (93.0)	
Yes	16 (5.7)	4 (7.0)	1.3 (0.5 to 3.5)
Intellectual ability			
Mean (SD)	1.5 (1.2)	3.2 (1.1)	3.2 (2.4 to 4.3)***
IQ≥50 (score 0, 1 or 2)	205 (73.5)	10 (17.5)	
IQ<50 (score 3 or 4)	72 (25.8)	47 (82.5)	
Missing	2 (0.7)	0	
Severity of motor disorder			
Hand manipulation (best hand) (0–4, 4 worst)			
Mean (SD)	1.5 (1.1)	3.3 (1.0)	
Score 0, 1 or 2	218 (78.1)	7 (12.3)	
Score 3 or 4	57 (20.4)	49 (86.0)	3.7 (2.7 to 5.0)***
Missing	4 (1.4)	1 (1.8)	
Able to sit by 24 months [†]			
Yes	131 (47.0)	1 (1.8)	
No	144 (51.6)	56 (98.3)	44.4 (6.1 to 320.8)***
Missing	4 (1.4)	0	
Able to cruise by 24 months [‡]			
Yes	102 (36.6)	2 (3.5)	
No	176 (63.1)	55 (96.5)	14.4 (3.5 to 59.2)***
Missing	1 (0.4)	0	

Continued

Table 1 Continued

	Alive (n=279)	Dead (n=57)	Hazard ratio (95% CI)
	No (%)	No (%)	
Disability predominantly in:			
Upper limb	14 (5.0)	2 (3.5)	
Lower limb	131 (47.0)	2 (3.5)	0.04 (0.0 to 0.2)
Equal	125 (44.8)	51 (89.5)	0.4 (0.1 to 1.6)***
Missing	9 (3.2)	2 (3.5)	
Type of CP predominantly:			
Spastic	250 (89.6)	50 (87.7)	
Ataxic	10 (3.6)	0	
Dyskinetic	12 (4.3)	6 (10.5)	2.3 (1.0 to 5.4)
Mixed	7 (2.5)	1 (1.8)	0.7 (0.1 to 4.9)
Bulbar involvement			
No	131 (47.0)	49 (85.0)	
Yes	136 (48.8)	7 (12.3)	6.3 (2.8 to 13.9)***
Missing	12 (4.3)	1 (1.8)	
Gastrostomy			
No	275 (98.6)	49 (86.0)	
Yes	4 (1.4)	7 (12.3)	5.1 (2.3 to 11.3)***
Fixed spinal curve			
No	265 (95.0)	53 (93.0)	
Yes	1 (0.4)	3 (5.3)	8.3 (2.6 to 26.6)***
Missing	13 (4.7)	1 (1.8)	
GMFCS score [§]			
I walks without limitations	39 (14.0)	0	
II walks with limitations	85 (30.5)	2 (3.5)	
III walks using a hand-held mobility device	34 (12.2)	0	
IV and V self-mobility with limitations, may use powered mobility; level V—transported in a manual wheelchair	121 (43.3)	55 (96.5)	6.2 (2.8 to 14.0)
Epilepsy			
Seizures in the last year			
None	193 (69.2)	14 (24.6)	
1–2	29 (10.4)	6 (10.5)	
3–6	23 (8.2)	3 (5.3)	
>6	24 (8.6)	30 (52.6)	2.2 (1.8 to 2.7)***
Missing	10 (3.6)	4 (7.0)	

*p<0.05, **p<0.01, ***p<0.001.

¹Getting to any sitting position from any lying position on the floor and then sitting without propping with either arm for 15 seconds.

²Cruising along furniture or a wall, even if placed in standing, for at least two steps sideways in either direction (not necessarily both).

[§]The Gross Motor Function Classification System (GMFCS)⁸ retrospectively assigned at age 2 to allow comparability to current studies. Levels IV and V combined owing to insufficient data at the age of 2 years to confidently distinguish them.

than 28 days postnatally (excluded by Hutton *et al*).¹ Interaction tests were carried out between these two factors and the other univariate factors.

RESULTS

Mortality

In all, 346 children were initially identified as alive from 1 year of age with bilateral CP (1.7/1000 live births). Only six (1.73%) could not be traced in 2006–2008. Sixty-one of the 340 children traced had died between the ages of 1 and 18 years—a rate of 17.94%.

The primary recorded cause of death in the majority was reported as respiratory: bronchopneumonia (19); respiratory infection (7); aspiration (6) (certificates for two of these cited gastro-oesophageal reflux contributing to aspiration and death); respiratory arrest (2) or respiratory failure (5). Eight died with epileptic seizure listed as the primary cause. The other primary causes were pancreatitis (1); diabetic ketoacidosis (1); hepatic failure (1); renal failure (1) and four gastrointestinal problems (three peritonitis and septicaemia associated with complications with gastrostomy/PEG and one bowel

infarction). In six cases, CP was given as the only cause of death, at ages ranging from 18 months to 13 years.

The survival of the cohort from the age of 1 year to 16–18 years is shown in figure 1. The mean age of death was 8.1 years (SD 5.2), but there were deaths throughout the period.

Late onset cases show a non-significant increased mortality over early onset (p=0.16). The mortality for children with a postnatal cause is also not significantly different from those with presumed prenatal and perinatal causation (p=0.58).

Predictors of mortality

Four of the 61 children died before 18 months, thus analyses of predictive factors from 2 years ± 6 months are based on 336 children (57 deaths and 279 surviving children).

Greater birth weight and a longer period of gestation were significantly associated with poorer survival (table 1). Poorer survival was also significantly associated with greater severity of motor disorder, as classified by the Gross Motor Function Classification System (GMFCS) or characterised by delayed acquisition of motor milestones, feeding and swallowing problems (as measured by the presence of a gastrostomy) and

Short report

severe communication problems. All but two deaths were in the combined level IV and V group. These exceptions were due, respectively, to diabetic ketoacidosis and epileptic seizure.

Visual impairment, epilepsy (more than six seizures in a year) and poor intellectual ability were non-motor measures of severity of disability significantly associated with poorer survival. Factors examined which were not significantly associated with survival were gender, clinical type of CP and respiratory infections indexed by reported wheeze and cough frequency at age 2 years.

The conclusions from these univariate analyses were not changed when taking into account age of onset (interaction tests $p > 0.05$).

The results from the multivariate analysis suggest that survival can be predicted by severely impaired hand function and/or communicative ability, and fixed spinal curvature. Removing the four participants affected by fixed spinal curvature at age 2 years from the analysis did not change this conclusion. Following Hutton *et al*,¹ we recoded both communication and hand manipulation problems into severe (scores 3 or 4) versus not severe (scores 0, 1 or 2). The hazard ratios and 95% CI for severe communication problems and for severe hand manipulation problems were highly significant (6.8 (1.9 to 23.9) and 5.7 (2.1 to 15.0), respectively).

Again following Hutton and colleagues,¹ we recoded these predictors into four categories: (1) severe communication *and* hand manipulation problems ($n=98$); (2) a severe communication problem but not a severe hand manipulation problem ($n=35$); (3) a severe hand manipulation problem but not a severe communication problem ($n=7$); or (4) *neither* a severe communication problem *nor* a severe hand manipulation problem ($n=189$). As expected, the survival was highest in those in category (4) and lowest in category (1) (figure 2). Of those children who had both severe manipulation and communication impairment, 42.7% survived to 18 years. However, when only one of either manipulation or communication function was severely impaired, there was 88% survival to 18 years.

DISCUSSION

This study has reported mortality from a complete population cohort of children with bilateral CP, who were all examined by one experienced clinician. Based on children who were alive at 1 year of age, the study had 98% ascertainment of deaths to age 16–18 years; 82% were still alive at age 16–18 years.




Figure 1 Survival of the cohort from the age of 1 year.

Important findings are that there is a steady mortality throughout childhood and adolescence; and that death is associated with severity of impairment from inferred bilateral brain damage affecting particularly the upper limbs and the bulbar region, with respiratory causes of death being most common.

Management of CP has changed over time, principally the use of gastrostomy in those with severe feeding problems and reflux. Strauss and colleagues have shown^{5,6} that survival for children who are largely immobile and fed by others (ie, the severely disabled group) has improved, possibly due to earlier gastrostomy. The present cohort study cannot address this point.

Brain insult may also give rise to cortical visual impairment and epilepsy. Both severe visual impairment and epilepsy were univariately associated with increased mortality in this study and a seizure was recorded as the primary cause of death in eight children. Spinal curvature already fixed at age 2 years, a likely indicator of severity of motor involvement, was also a risk factor. Hearing loss, thought at one time to be an additional mortality risk factor, was not a significant risk factor in this study.

The mortality we report is likely to be an underestimate of the birth cohort mortality of children with bilateral CP, as children who died before 12 months of age were never seen to confirm diagnosis and thus never included in the study. Variation in inclusion criteria may have an impact on results. This cohort mortality is a little higher than that in the study by Hutton *et al*,¹ which described mortality in a birth cohort of CP registered from 1 month of age (whereas our study comprised those alive at 12 months; excluded onset after 28 days post birth; and had a mix of unilateral and bilateral CP), and where the 20-year survival was 89.3% for females and 86.9% for males.

Of the other factors noted previously, we found that the hazard ratio for female gender was also reduced,¹ although this reduction did not reach conventional statistical significance; and that larger birth weight and greater gestational age were also factors relevant to survival,¹ although less predictive than functional impairment. This latter could be due to a greater infant death rate in those born prematurely and of low birth weight. Another factor may be that those with a postneonatal brain insult have increased mortality (and greater functional impairment) but are more likely to have been of larger birth weight and greater gestational age.

We have shown that easily ascertainable measures are relevant to prognosis and are presented here in a way that enables




Figure 2 Survival of cohort from 18 months split by presence/absence of communication and manipulation problems.

their use in conversations with parents. In children with bilateral CP, parents and clinicians can confidently expect most children with bilateral CP who at the age of 2 years have good or mildly impaired communication and manipulation skills with no bulbar problems, and who can cruise and can get to sit, will live to 18 years and—according to data from UK and USA studies⁷—well beyond.

Acknowledgements The authors would like to thank all the families, especially the young people, for their participation; also Alison Davis for data collection, Felicity Clemens for early statistical advice and Lewis Rosenbloom for helpful suggestions.

Funding The study was funded by the Charles Wolfson Charitable Trust, Cerebra, One Small Step and the Hickman Fund (both these last administered by Ruth Bishop, Special Purpose Funds Manager, Guy's and St Thomas' Charity).

Competing interests None.

Ethics approval Ethical approval was granted by the South-East MREC, reference number 06/MRE01/18.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. **Hutton JL**, Cooke T, Pharoah PO. Life expectancy in children with cerebral palsy. *BMJ* 1994;**309**:431–5.
2. **Strauss DJ**, Shavelle RM, Anderson TW. Life expectancy of children with cerebral palsy. *Pediatr Neurol* 1998;**18**:143–9.
3. **Blair E**, Watson L, Badawi N, *et al*. Life expectancy among people with cerebral palsy in Western Australia. *Dev Med Child Neurol* 2001;**43**:508–15.
4. **Scrutton D**, Baird G, Smeeton N. Hip dysplasia in bilateral cerebral palsy: incidence and natural history in children aged 18 months to 5 years. *Dev Med Child Neurol* 2001;**43**:586–600.
5. **Strauss D**, Shavelle R, Reynolds R, *et al*. Survival in cerebral palsy in the last 20 years: signs of improvement? *Dev Med Child Neurol* 2007;**49**:86–92.
6. **Strauss D**, Brooks J, Rosenbloom L, *et al*. Life expectancy in cerebral palsy: an update. *Dev Med Child Neurol* 2008;**50**:487–93.
7. **Hemming K**, Hutton JL, Pharoah PO. Long-term survival for a cohort of adults with cerebral palsy. *Dev Med Child Neurol* 2006;**48**:90–5.
8. **GMFCS—Expanded and Revised (2007)**. <http://motorgrowth.canchild.ca/en/GMFCS/expandedandrevised.asp> (Accessed 12 January 2010).



Mortality from 1 to 16–18 years in bilateral cerebral palsy

Gillian Baird, Elizabeth Allen, David Scrutton, et al.

Arch Dis Child 2011 96: 1077-1081 originally published online June 15, 2010

doi: 10.1136/adc.2009.172841

Updated information and services can be found at:

<http://adc.bmj.com/content/96/11/1077.full.html>

These include:

References

This article cites 7 articles, 1 of which can be accessed free at:

<http://adc.bmj.com/content/96/11/1077.full.html#ref-list-1>

Article cited in:

<http://adc.bmj.com/content/96/11/1077.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Epidemiologic studies](#) (1047 articles)
[Child health](#) (2259 articles)
[Cerebral palsy](#) (82 articles)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>